

A ROBIN FORMULA FOR THE FEKETE-LEJA TRANSFINITE DIAMETER

ROBERT RUMELY

The purpose of this note is to give a formula for the Fekete-Leja transfinite diameter on $\mathbb{C}^N$, generalizing the classical Robin formula

$$d_\infty(E) = e^{-V(E)}$$

for the usual transfinite diameter.

We will disengage this from a formula for the sectional capacity proved in arithmetic intersection theory ([5], Theorem 1.1, p.233).

First recall the definition of the Fekete-Leja transfinite diameter for a compact set $E \subset \mathbb{C}^N$ (see [1], [16]). Consider the set of monomials $z^k = z_1^{k_1} \cdots z_N^{k_N}$ in the polynomial ring $\mathbb{C}[z] = \mathbb{C}[z_1, \dots, z_N]$. Let $\Gamma(n) \subset \mathbb{C}[z]$ be the space of polynomials of total degree at most n , and let $\mathcal{K}(n) = \{k \in \mathbb{Z}^N : k_i \geq 0, k_1 + \cdots + k_N \leq n\}$ be the index set for the monomial basis of $\Gamma(n)$. Put $q_n = \#(\mathcal{K}(n)) = \binom{n+N}{N}$.

Fixing n , take q_n independent vector variables $z_i = (z_{i1}, \dots, z_{iN}) \in \mathbb{C}^N$, $i = 1, \dots, q_n$. Let $k_1, \dots, k_{q_n}$ be the indices in $\mathcal{K}(n)$. The Vandermonde determinant

$$Q_n(z_1, \dots, z_{q_n}) := \det(z_i^{k_j})_{i,j=1}^{q_n}$$

is a homogeneous polynomial in the z_{ij} of total degree $T_n = N \cdot \binom{n+N}{N+1}$. For each n , put

$$d_n(E) = \max_{z_1, \dots, z_{q_n} \in E} |Q_n(z_1, \dots, z_{q_n})|^{1/T_n}.$$

The Fekete-Leja transfinite diameter is defined by

$$d_\infty(E) = \lim_{n \rightarrow \infty} d_n(E).$$

The existence of the limit is due to Zaharjuta ([16]).

Henceforth, we will assume that $d_\infty(E) > 0$. This is equivalent to E being non-pluripolar ([10]).

For $f \in \mathbb{C}[z_1, \dots, z_N]$ write $\|f\|_E = \sup_{z \in E} |f(z)|$. The Green's function $G^*(z, E)$ is the upper semicontinuous regularization of the Siciak extremal function

$$G(z, E) := \lim_{n \rightarrow \infty} \max_{\substack{f \in \Gamma(n) \\ \|f\|_E = 1}} \frac{1}{n} \log(|f(z)|).$$

Date: September 21, 2005.

2000 Mathematics Subject Classification. Primary: 32U35, 32U20, 14G40; Secondary: 31B15, 31C10.

Key words and phrases. Fekete-Leja transfinite diameter, Robin formula.

Work supported in part by NSF grant DMS-0300784.

Since E is not pluripolar, $G^*(z, E)$ is finite for all $z \in \mathbb{C}^N$ and is plurisubharmonic ([9], [10], [17]). Write $dd^c = \frac{i}{4\pi} \partial \bar{\partial}$ and let

$$(1) \quad \omega = dd^c(2G^*(z, E))$$

be the associated positive $(1, 1)$ -current. (The factor 2 is needed for compatibility with the Poincaré-Lelong formula).

Let $Z_0, \dots, Z_N$ be homogeneous coordinates on $\mathbb{P}^N(\mathbb{C})$. For $k = 0, \dots, N$ write H_k for the hyperplane $\{Z_k = 0\}$ and let $\mathbb{A}_k^N = \mathbb{P}^N(\mathbb{C}) \setminus H_k$ be the corresponding affine patch. Identify $\mathbb{C}^N$ with $\mathbb{A}_0^N \subset \mathbb{P}^N(\mathbb{C})$ via the embedding $(z_1, \dots, z_N) \mapsto (1 : z_1 : \dots : z_N)$. If $\|z\| = (|z_1|^2 + \dots + |z_N|^2)^{1/2}$, then $|G^*(z, E) - \max(0, \log(\|z\|))|$ is uniformly bounded on $\mathbb{C}^N$ (for boundedness from above, see [9], Corollary 5.2.2, p.193; for boundedness from below, note that since E is compact, it is contained in a ball $B(0, R) = \{z \in \mathbb{C}^N : \|z\| \leq R\}$ for some large R , so $G^*(z, E) \geq G^*(z, B(0, R)) = \max(0, \log(\|z\|/R))$). It follows that for each $1 \leq k \leq N$ the function $g_k(z, E)$ defined on $\mathbb{A}_k^N \setminus H_0$ by

$$g_k(z, E) = G^*(z, E) - \log(|z_k|)$$

extends uniquely to a plurisubharmonic function on $\mathbb{A}_k^N$. If $k = 0$, write $g_0(z, E) = G^*(z, E)$. Since $\log(|Z_i/Z_j|)$ is pluriharmonic on $\mathbb{A}_i^N \cap \mathbb{A}_j^N$, the currents $\omega_k := dd^c(2g_k(z, E))$ cohere to give a positive $(1, 1)$ -current on $\mathbb{P}^N(\mathbb{C})$ extending ω . We will denote it by ω as well.

Let $\omega^k = \omega \wedge \dots \wedge \omega$ be the k -fold exterior product of ω with itself, and put $\omega^0 = 1$. Write Y_k for the space $H_0 \cap \dots \cap H_{k-1}$ and let $U_k = Y_k \setminus H_k$, so $U_k(\mathbb{C}) \cong \mathbb{C}^{N-k}$. Define the *iterated Robin constant* by

$$(2) \quad \begin{aligned} \tilde{V}(E) &= \frac{1}{N} \sum_{k=1}^N \int_{U_k(\mathbb{C})} g_k(z, E) \omega^{N-k} \\ &= \frac{1}{N} \left(\int_{Z_0=0, Z_1 \neq 0} g_1(z, E) \omega^{N-1} \right. \\ (3) \quad &\quad \left. + \int_{Z_0=Z_1=0, Z_2 \neq 0} g_2(z, E) \omega^{N-2} \right. \\ &\quad \left. + \dots + g_N((0 : \dots : 0 : 1), E) \right). \end{aligned}$$

Note that when $N = 1$, the sum consists of a single term and reduces to the usual Robin constant, since $\infty = (0 : 1) \in \mathbb{P}^1(\mathbb{C})$, and $g_1(\infty, E) = \lim_{z \rightarrow \infty} G^*(z, E) - \log(|z|) = V(E)$.

Theorem 0.1. *If $E \subset \mathbb{C}^N$ is compact and not pluripolar, then*

$$(4) \quad d_\infty(E) = e^{-\tilde{V}(E)}.$$

Before giving the proof we will need some facts from arithmetic intersection theory. Let K be a number field, and let $\mathcal{O}_K$ be the ring of integers of K . Let X/K be a smooth, connected projective variety of dimension N , and write $K(X)$ for the field of K -rational functions on X .

The sectional capacity $S_\gamma(\mathbb{E}, D)$ is a measure of size for an adelic set $\mathbb{E}$ on X relative to an ample divisor D . It was first proposed by Chinburg ([4]), and its existence was shown in ([14]), using methods from ([16]). The name ‘sectional capacity’ refers to asymptotics of volumes of spaces of sections of $\mathcal{O}_X(nD)$ as $n \rightarrow \infty$.

Let D be an effective, ample K -rational Cartier divisor on X .

For each place v of K , let K_v be the completion of K at v , and let $\mathbb{C}_v$ be the minimal complete algebraically closed field containing K_v . Let $|x|_v$ be the absolute value on $\mathbb{C}_v$ extending the canonical absolute value on K_v given by the modulus of additive Haar measure. Without loss, we can assume that X is embedded in $\mathbb{P}^M$ for a suitable M . There is a natural distance function $d_v(x, y)$ on $\mathbb{P}^M(\mathbb{C}_v)$: the chordal metric associated to the Fubini-Study metric, if v is archimedean; the v -adic spherical metric, if v is nonarchimedean (see [12], §1.1).

For each v , let $E_v \subset X(\mathbb{C}_v)$ be a nonempty set, and write $\mathbb{E} = \prod_v E_v$; we will call $\mathbb{E}$ an *adelic set*. We will assume that the E_v and D satisfy the following ‘Standard Hypotheses’:

(1) Each E_v is bounded away from $\text{supp}(D)(\mathbb{C}_v)$ under $d_v(x, y)$ and is stable under the group of continuous automorphisms $\text{Gal}^c(\mathbb{C}_v/K_v)$.

(2) For all but finitely many v , E_v and $\text{supp}(D)(\mathbb{C}_v)$ specialize to disjoint sets (mod v); equivalently, for all but finitely many v , $d_v(x, y) = 1$ for all $x \in E_v$ and all $y \in \text{supp}(D)(\mathbb{C}_v)$.

Note that if v is archimedean and $K_v \cong \mathbb{R}$, then $\text{Gal}^c(\mathbb{C}_v/K_v) = \{1, \tau\}$ where τ is complex conjugation; if $K_v \cong \mathbb{C}$, then $\text{Gal}^c(\mathbb{C}_v/K_v)$ is trivial.

For each integer $n \geq 0$, put $\Gamma(nD) = H^0(X, \mathcal{O}_X(nD))$, a finite dimensional vector space over K . Dehomogenizing at D , identify $\Gamma(nD)$ with $\{f \in K(X) : \text{div}(f) + nD \geq 0\}$. For each v , consider the space of K_v -rational functions on X with polar divisor at most nD , $\Gamma_v(nD) = K_v \otimes_K \Gamma(nD) = \{f \in K_v(X) : \text{div}(f) + nD \geq 0\}$. For $f \in K_v(X)$, write $\|f\|_{E_v} = \sup_{x \in E_v} |f(x)|_v$ and let

$$\mathcal{F}(E_v, nD) = \{f \in \Gamma_v(nD) : \|f\|_{E_v} \leq 1\}.$$

Let $K_{\mathbb{A}} = \{(x_v) \in \prod_v K_v : |x_v|_v \leq 1 \text{ for all but finitely many } v\}$ be the adèle ring of K . Identifying $K_{\mathbb{A}} \otimes_K \Gamma(nD)$ with a subset of $(\prod_v K_v) \otimes_K \Gamma(nD) \cong \prod_v (K_v \otimes_K \Gamma(nD))$, introduce the ‘adelic unit ball’

$$\mathcal{F}(\mathbb{E}, nD) = (K_{\mathbb{A}} \otimes_K \Gamma(nD)) \bigcap \left(\prod_v \mathcal{F}(E_v, nD) \right).$$

Fix an additive Haar measure $\text{vol}_{\mathbb{A}}$ on $K_{\mathbb{A}}$. By transport of structure using a K -basis for $\Gamma(nD)$, $\text{vol}_{\mathbb{A}}$ induces a Haar measure on $K_{\mathbb{A}} \otimes_K \Gamma(nD)$. The Product Formula shows this measure is independent of the choice of basis; it too will be denoted $\text{vol}_{\mathbb{A}}$. The sectional capacity $S_{\gamma}(\mathbb{E}, D)$ is defined by

$$-\log(S_{\gamma}(\mathbb{E}, D)) = \lim_{n \rightarrow \infty} \frac{(N+1)!}{n^{N+1}} \text{vol}_{\mathbb{A}}(\mathcal{F}(\mathbb{E}, nD))$$

(see [14], pp.23-24). The limit is independent of the choice of Haar measure on $K_{\mathbb{A}}$. Its existence is part of ([14], Theorem C, p.8).

We will now discuss Arakelov theory. Let $\mathfrak{X}$ be a model of X : an integral projective scheme of Krull dimension $N+1$, flat and proper over $\text{Spec}(\mathcal{O}_K)$, whose generic fibre is K -isomorphic to X .

Put $X_{\mathbb{C}} = X \times_{\mathbb{Q}} \text{Spec}(\mathbb{C})$; then $X_{\mathbb{C}}(\mathbb{C}) = \bigsqcup_{\sigma} X_{\sigma}(\mathbb{C})$ is a complex manifold with a component corresponding to each embedding $\sigma : K \hookrightarrow \mathbb{C}$. Write τ for complex conjugation on $\mathbb{C}/\mathbb{R}$; there is a natural operation of τ on $X_{\mathbb{C}}(\mathbb{C})$ coming from its action on the base. If σ is a real embedding, then τ acts as complex conjugation on $X_{\sigma}(\mathbb{C})$; if $\{\sigma, \bar{\sigma}\}$ is a pair of conjugate complex embeddings, then τ interchanges the components X_{σ} and $X_{\bar{\sigma}}$, mapping $P \in X_{\sigma}(\mathbb{C})$

to $\overline{P} \in X_{\overline{\sigma}}(\mathbb{C})$, where $\overline{P}$ is the point whose coordinates are the complex conjugates of those of P .

Let L be a line bundle on X . A *metrized line bundle* $\overline{\mathcal{L}} = (\mathcal{L}, [\cdot])$ on $\mathfrak{X}$, extending L , is a pair consisting of a locally free sheaf $\mathcal{L}$ of rank 1 on $\mathfrak{X}$ which induces L on X , and a smooth positive Hermitian metric $[\cdot]$ on the fibres of L over $X_{\mathbb{C}}(\mathbb{C})$, which is invariant under the action of τ . Let $c_1(\overline{\mathcal{L}})$ be the first Chern class of $\overline{\mathcal{L}}$, the smooth $(1, 1)$ -form on $X_{\mathbb{C}}(\mathbb{C})$ defined in a neighborhood of any point $z_0 \in X_{\mathbb{C}}(\mathbb{C})$ by $dd^c(-2\log([\cdot](s(z))))$, where s is a meromorphic section of L which is defined and does not vanish at z_0 .

For each cycle $\mathcal{Y}$ on $\mathfrak{X}$, Bost, Gillet and Soulé ([2]) define the “height” $h_{\overline{\mathcal{L}}}(\mathcal{Y})$ of $\mathcal{Y}$ relative to $\overline{\mathcal{L}}$ to be the self-intersection number $\widehat{\deg}(\hat{c}_1(\overline{\mathcal{L}})^{\dim(\mathcal{Y})}|_{\mathcal{Y}})$ in the arithmetic intersection theory Gillet and Soulé developed in ([6], [7], [8]). The height is additive in $\mathcal{Y}$. There is a recursive formula for $h_{\overline{\mathcal{L}}}(\mathcal{Y})$ (see [2], Proposition 3.2.1(iv), p.949, and the remarks after it dealing with the non-regular case): when $\mathcal{Y}$ is irreducible, let $s \neq 0$ be a section of $\mathcal{L}|_{\mathcal{Y}}$, and let $\text{div}_{\mathcal{Y}}(s)$ be its divisor. Write Y for the generic fibre $\mathcal{Y}_K$. Then

$$(5) \quad h_{\overline{\mathcal{L}}}(\mathcal{Y}) = \int_{Y_{\mathbb{C}}(\mathbb{C})} -\log([\cdot](s(z))) c_1(\overline{\mathcal{L}})^{\dim(Y)} + h_{\overline{\mathcal{L}}}(\text{div}_{\mathcal{Y}}(s)).$$

Eventually the recursion abuts at a 0-cycle $\mathcal{Z} = \sum_{\mathfrak{p}} n_{\mathfrak{p}} \cdot \mathfrak{p}$, a finite sum of closed points, and then $h_{\overline{\mathcal{L}}}(\mathcal{Z}) = \sum n_{\mathfrak{p}} \log(N_{\mathfrak{p}})$ where $N_{\mathfrak{p}}$ is the order of the residue field at $\mathfrak{p}$ ([2], formula 3.1.4, p.946).

It is customary to write $\overline{\mathcal{L}}^{N+1}$ for $h_{\overline{\mathcal{L}}}(\mathfrak{X})$. If $\overline{\mathcal{L}}$ is replaced by $\overline{\mathcal{L}}^{\otimes m}$, then $(\overline{\mathcal{L}}^{\otimes m})^{N+1} = m^{N+1} \cdot \overline{\mathcal{L}}^{N+1}$ ([2], Proposition 3.2.1(i)). This leads to the notion of a *fractional metrized line bundle*: if $n > 0$ is an integer, and $\overline{\mathcal{L}}$ is a metrized line bundle inducing $L^{\otimes n}$ on X , then we call the formal object $\frac{1}{n}\overline{\mathcal{L}}$ a fractional metrized line bundle, and define $(\frac{1}{n}\overline{\mathcal{L}})^{N+1} = n^{-(N+1)} \cdot \overline{\mathcal{L}}^{N+1}$.

Chinburg, Lau and Rumely ([5]) expressed the sectional capacity as a limit of self-intersection numbers of fractional metrized line bundles. Given an adelic set $\mathbb{E}$ and an effective ample divisor D on X satisfying the Standard Hypotheses, they constructed a sequence of models $\mathfrak{X}_n$, and metrized line bundles $\overline{\mathcal{L}}_n$ on $\mathfrak{X}_n$ extending $L^{\otimes n} = \mathcal{O}_X(nD)$ on X , such that

$$(6) \quad -\log(S_{\gamma}(\mathbb{E}, D)) = \lim_{n \rightarrow \infty} \left(\frac{1}{n} \overline{\mathcal{L}}_n \right)^{N+1} :$$

see ([5], Theorem 1.1, p.233).

The models $\mathfrak{X}_n$ are defined using the nonarchimedean part of $\mathbb{E}$. For each n , put $S_n = \{f \in \Gamma(nD) : \|f\|_{E_v} \leq 1 \text{ for all nonarchimedean } v\}$, and let $\mathcal{O}_K[S_n]$ be the graded $\mathcal{O}_K$ -algebra generated in degree 1 by S_n . Then $\mathfrak{X}_n = \text{Proj}(\mathcal{O}_K[S_n])$. Let $\mathcal{L}_n = \mathcal{O}_{\mathfrak{X}_n}(1)$; then $\mathcal{L}_n$ induces $L^{\otimes n}$ on X .

The metrics $[\cdot]_n$ are constructed using the archimedean part of $\mathbb{E}$. For each archimedean place v of K , fix an isomorphism $\mathbb{C}_v \cong \mathbb{C}$, and choose a sequence of sets $E_{v,1} \supseteq E_{v,2} \supseteq \cdots$ containing E_v for which the extremal functions

$$G(z, E_{v,n}, D) := \lim_{m \rightarrow \infty} \sup_{\substack{f \in \Gamma(mD) \\ \|f\|_{E_{v,n}} \leq 1}} \frac{1}{m} \log(|f(z)|)$$

are continuous and increase monotonically to $G(z, E_v, D)$. (The existence of such sets $E_{v,n}$ follows from the proof of ([5], Lemma 1.2, p.234).)

Since $G(z, E_{v,n}, D)$ is continuous, it is plurisubharmonic. By a theorem of Richburg ([11], Satz 4.7) there is a smooth plurisubharmonic function $G_{v,n}(z)$ such that

$$G(z, E_{v,n}, D) - \frac{1}{n} \leq G_{v,n}(z) \leq G(z, E_{v,n}, D) - \frac{1}{n+1}$$

for all $z \in X(\mathbb{C}_v) \setminus \text{supp}(D)(\mathbb{C}_v)$. Hence

$$G(z, E_v, D) = \lim_{n \rightarrow \infty} G_{v,n}(z)$$

as an increasing limit.

By ([5], Theorem 2.13, p.253), the smoothings $G_{v,n}(z)$ can be chosen so that for each $z_0 \in \text{supp}(D)(\mathbb{C}_v)$, if s is a local equation for D at z_0 , then $G_{v,n}(z) + \log(|s(z)|)$ extends to a smooth plurisubharmonic function in a neighborhood of z_0 . If $K_v \cong \mathbb{R}$, then since E_v has been assumed to be stable under complex conjugation, the $G_{v,n}(z)$ can be chosen to be invariant under complex conjugation as well.

Each embedding $\sigma : K \hookrightarrow \mathbb{C}$ determines a place v of K . If σ is a real embedding, put $G_{\sigma,n}(z) = G_{v,n}(z)$. If σ is a complex embedding, then precisely one of σ and its complex conjugate $\bar{\sigma}$ induces the chosen isomorphism $\mathbb{C}_v \cong \mathbb{C}$; if it is σ , put $G_{\sigma,n}(z) = G_{v,n}(z)$, if not, put $G_{\sigma,n}(z) = G_{v,n}(\bar{z})$.

The metric $\llbracket \cdot \rrbracket_n$ is defined by requiring that for the tautological section 1 of $L^{\otimes n}$, if $z \in X_{\sigma}(\mathbb{C}) \subset X_{\mathbb{C}}(\mathbb{C})$ then

$$\llbracket 1(z) \rrbracket_n = \exp(-nG_{\sigma,n}(z)) .$$

By construction, $\llbracket \cdot \rrbracket_n$ is invariant under τ .

Proof of Theorem 0.1. Let $E \subset \mathbb{C}^N$ be a compact, non-pluripolar set. We will embed it as the archimedean component of an adelic set, and apply the machinery above.

Take $K = \mathbb{Q}(\sqrt{-1})$ as the ground field, so $\mathcal{O}_K$ is the ring of Gaussian integers. Write $\mathbb{P}_{\mathcal{O}_K}^N$ for $\mathbb{P}^N/\text{Spec}(\mathcal{O}_K)$, and write $\mathbb{P}_K^N$ for its generic fibre. Let X/K be $\mathbb{P}_K^N$, and let D be defined by $\{Z_0 = 0\}$.

Put $\mathbb{A}^N = \mathbb{P}_K^N \setminus H_0$, and define $\mathbb{E} = \prod_v E_v \subset \prod_v \mathbb{A}^N(\mathbb{C}_v)$ as follows. For each place v , identify $\mathbb{A}^N(\mathbb{C}_v)$ with $\mathbb{C}_v^N$. There is one archimedean place v_{∞} for K ; fix an isomorphism $K_{v_{\infty}} \cong \mathbb{C}$ and put $E_{v_{\infty}} = E \subset \mathbb{C}^N$. Condition (1) of the Standard Hypotheses holds trivially. For each nonarchimedean v , put $E_v = B(0, 1) = \{(z_1, \dots, z_N) \in \mathbb{C}_v^N : \max(|z_i|_v) \leq 1\}$. Since $\text{Gal}^c(\mathbb{C}_v/K_v)$ preserves $|x|_v$, again condition (1) of the Standard Hypotheses holds. The sets E_v and $H_0(\mathbb{C}_v)$ specialize to disjoint sets (mod v) for all nonarchimedean v , so condition (2) of the Standard Hypotheses holds as well.

By ([13], Theorem 3.1, p.551) the sectional capacity $S_{\gamma}(\mathbb{E}, D)$ can be decomposed as a product of ‘local sectional capacities’

$$S_{\gamma}(\mathbb{E}, D) = \prod_v S_{\gamma}(E_v, D) .$$

Here the $S_{\gamma}(E_v, D)$ depend on the choice of an ordered basis for the graded ring $\bigoplus_{n=0}^{\infty} \Gamma(nD)$: we take this to be the monomial basis, equipped with the lexicographic order graded by the degree. In this situation, $S_{\gamma}(E_{v_{\infty}}, D) = d_{\infty}(E)^{2N}$. This follows from ([13], Theorems 2.3 and 2.6), combined with the discussion on ([13], p.557). (In ([13], p.557), the formula $S_{\gamma}(E_{\infty}, D) = d_{\infty}(E)^N$ is given when $\mathbb{R}$ is the ground field. Here, since $K_{v_{\infty}} \cong \mathbb{C}$, the normalized absolute value $|x|_{v_{\infty}} = |x|^2$ used in computing $S_{\gamma}(E_{v_{\infty}}, D)$ is the square of the

usual absolute value; this is the source of the 2 in the exponent.) Furthermore, $S_\gamma(E_v, D) = 1$ for each nonarchimedean v by ([13], Example 4.1, p.555). Hence

$$(7) \quad S_\gamma(\mathbb{E}, D) = d_\infty(E)^{2N}.$$

Let $\{\frac{1}{n}\overline{\mathcal{L}}_n\}$ be the sequence of fractional metrized line bundles constructed in ([5], Theorem 1.1) as described above. Since E_v is the ‘trivial set’ $B(0, 1)$ for each nonarchimedean v , the Maximum Modulus Principle of nonarchimedean analysis shows that

$$S_n = \oplus_{k_0+\dots+k_N=n} \mathcal{O}_K \cdot Z_0^{k_0} \cdots Z_N^{k_N}.$$

It follows that $\mathfrak{X}_n \cong \mathbb{P}_{\mathcal{O}_K}^N$, and $\mathcal{L}_n \cong \mathcal{O}_{\mathbb{P}_{\mathcal{O}_K}^N}(n)$. Let $\overline{\mathcal{L}}'_n$ be $\mathcal{O}_{\mathbb{P}_{\mathcal{O}_K}^N}(1)$, equipped with the metric $\llbracket \cdot \rrbracket'_n$ defined by $\llbracket 1(z) \rrbracket'_n = \exp(-G_n(z))$, where $G_n(z)$ is metric constructed as above by smoothing $G(z, E, D)$. Then $\overline{\mathcal{L}}_n \cong (\overline{\mathcal{L}}'_n)^{\otimes n}$, so $(\frac{1}{n}\overline{\mathcal{L}}_n)^{N+1} = (\overline{\mathcal{L}}'_n)^{N+1}$ and without loss we can replace $\frac{1}{n}\overline{\mathcal{L}}_n$ by $\overline{\mathcal{L}}'_n$.

We will now compute the intersection product $(\overline{\mathcal{L}}'_n)^{N+1}$ using Bost-Gillet-Soulé’s recursive formula. There are two embeddings of K into $\mathbb{C}$, both of which correspond to the place v_∞ , so $(\mathbb{P}_K^N)_{\mathbb{C}}(\mathbb{C})$ has two components which are interchanged by τ . Write $\mathbb{P}^N(\mathbb{C})$ for the one corresponding to our chosen isomorphism $K_{v_\infty} \cong \mathbb{C}$, and $(\mathbb{P}^N)_\tau(\mathbb{C})$ for the other. Since they are isomorphic, and the metrics $\llbracket \cdot \rrbracket_n$ are τ -invariant, the integrals over the two components in the archimedean part of the intersection product are the same. Therefore, in what follows, we will compute the integrals over $\mathbb{P}^N(\mathbb{C})$ and double the answer. For any K -rational subvariety $Y \subset \mathbb{P}_K^N$, write $Y(\mathbb{C})$ for the part of $Y_{\mathbb{C}}(\mathbb{C})$ in the chosen component $\mathbb{P}^N(\mathbb{C})$.

Write $\mathbb{A}_k^N$ for $\mathbb{P}_K^N \setminus H_k$, where $H_k = \{Z_k = 0\}$. Let $\overline{H}_k$ be the Zariski closure of H_k in $\mathbb{P}_{\mathcal{O}_K}^N$. Then $\overline{H}_0, \dots, \overline{H}_N$ meet transversely on $\mathbb{P}_{\mathcal{O}_K}^N$. Write $g_{n,0}(x) = G_n(x)$ on $(\mathbb{A}_0^N)(\mathbb{C})$ and for each $k = 1, \dots, N$ let $g_{n,k}(x)$ be the natural extension of $G_n(x) - \log(|z_k(x)|)$ to a plurisubharmonic function on $\mathbb{A}_k^N(\mathbb{C})$; here $z_k = Z_k/Z_0$. As before, the $(1, 1)$ -forms $dd^c(2g_{n,k})$ glue to give a well-defined $(1, 1)$ -form ω_n on $\mathbb{P}^N(\mathbb{C})$.

Since $L = \mathcal{O}_{\mathbb{P}_K^N}(D)$ where $D = \{Z_0 = 0\}$, the canonical section ‘1’ of L is Z_0 . This means that for each $x \in \mathbb{A}_0^N(\mathbb{C})$

$$-\log(\llbracket Z_0(x) \rrbracket'_n) = G_n(x) = g_{n,0}(x).$$

Similarly, on $\mathbb{A}_k^N(\mathbb{C})$,

$$\begin{aligned} -\log(\llbracket Z_k(x) \rrbracket'_n) &= -\log(\llbracket (Z_k/Z_0)(x) \cdot Z_0(x) \rrbracket'_n) \\ &= -\log(|z_k(x)|) + G_n(x) = g_{n,k}(x). \end{aligned}$$

Put $\mathcal{Y}_0 = \mathfrak{X}_n = \mathbb{P}_{\mathbb{Z}}^N$ and take $s = Z_0$. Let $Y_0 = \mathbb{P}_{\mathbb{Q}}^N$ be the generic fibre of $\mathcal{Y}_0$ and put $\mathcal{Y}_1 = \text{div}(Z_0) = \overline{H}_0$. By Bost-Gillet-Soulé’s formula (5) and the remarks above,

$$(\overline{\mathcal{L}}'_n)^{N+1} = h_{\overline{\mathcal{L}}'_n}(\mathfrak{X}_n) = 2 \int_{Y_0(\mathbb{C})} g_{n,0}(z) \omega^N + h_{\overline{\mathcal{L}}'_n}(\mathcal{Y}_1).$$

Inductively apply this formula to the sections $Z_1, \dots, Z_N$, putting $\mathcal{Y}_{k+1} = \text{div}_{\mathcal{Y}_k}(Z_k|_{\mathcal{Y}_k}) = \overline{H}_0 \cdot \dots \cdot \overline{H}_k$. Note that $\overline{H}_0 \cdot \dots \cdot \overline{H}_N = 0$, so the final abutment term vanishes. It follows that

$$(8) \quad (\overline{\mathcal{L}}'_n)^{N+1} = 2 \sum_{k=0}^N \int_{Y_k(\mathbb{C})} g_{n,k}(z) \omega_n^{N-k}.$$

Since ω_n is smooth, the $(N - k - 1)$ -dimensional subspace $Y_{k+1}(\mathbb{C})$ of $Y_k(\mathbb{C})$ has measure 0 under ω_n^{N-k} . Put $U_k = Y_k \setminus Y_{k+1}$. Then for each k

$$(9) \quad \int_{Y_k(\mathbb{C})} g_{n,k}(z) \omega_n^{N-k} = \int_{U_k(\mathbb{C})} g_{n,k}(z) \omega_n^{N-k}.$$

Now let $n \rightarrow \infty$. For each k the $g_{n,k}$ increase monotonically on $U_k(\mathbb{C})$ to $G(z, E) - \log(|z_k|)$, whose upper semicontinuous regularization is $g_k(z, E)$. Noting that $U_k(\mathbb{C}) \cong \mathbb{C}^{N-k}$, it follows from ([3], Theorem 7.4) that

$$(10) \quad \lim_{n \rightarrow \infty} \int_{U_k(\mathbb{C})} g_{n,k}(z) \omega_n^{N-k} = \int_{U_k(\mathbb{C})} g_k(z, E) \omega^{N-k}.$$

Combining (6), (8), (9), and (10) gives

$$(11) \quad -\log(S_\gamma(\mathbb{E}, D)) = 2 \sum_{k=0}^N \int_{U_k(\mathbb{C})} g_k(z, E) \omega^{N-k}.$$

When $k = 0$, $g_0(z, E)$ is the extremal plurisubharmonic function $G^*(z, E)$ on $U_0(\mathbb{C}) \cong \mathbb{C}^N$, and ω^N is the Bedford-Taylor measure, which is supported on E . The set $\{z \in E : G^*(z, E) > 0\}$ is a negligible set, so it has measure 0 under ω^N . Hence

$$(12) \quad \int_{U_0(\mathbb{C})} g_0(z, E) \omega^N = 0.$$

Using (7), (11), and (12) we obtain

$$-2N \cdot \log(d_\infty(E)) = 2 \sum_{k=1}^N \int_{U_k(\mathbb{C})} g_k(z, E) \omega^{N-k}$$

which is equivalent to (4). □

Generalizations. It is tempting to assert that the formula in Theorem 0.1 gives a new definition of the capacity. However, we do not do so because the stated formula is only one of a class of formulas with the same property.

In particular, the order in which the hyperplanes H_k are intersected in the definition of $\tilde{V}(E)$ is immaterial. More generally, if $A = (a_{kj}) \in \text{GL}_N(\mathbb{C})$, and if $Z'_k = \sum a_{kj} Z_j$ for $k = 1, \dots, N$ then the constant $\tilde{V}(E)$ can equally well be defined by

$$(13) \quad \tilde{V}(E) = \frac{1}{N} \sum_{k=1}^N \int_{Z_0=Z'_1=\dots=Z'_{k-1}=0, Z'_k \neq 0} g'_k(z, E) \omega^{N-k} + \frac{1}{N} \log(|\det(A)|)$$

where $g'_k(z, E) = G^*(z, E) - \log(|\sum a_{kj} z_j|)$ on $\mathbb{C}^N$.

Indeed, if we put $z' = {}^t(z'_1, \dots, z'_N) = A(z)$, where $z'_k = \sum a_{kj} z_j$ for $k = 1, \dots, N$, this follows from Sheinov's formula (see [15], or [1], p.287)

$$d_\infty(A(E)) = |\det(A)|^{1/N} \cdot d_\infty(E),$$

by applying Theorem 0.1 to the set $A(E)$, and noting that $G(z', A(E)) = G(z, E)$, as follows easily from the definitions.

REFERENCES

- [1] T. Bloom and J.P. Calvi, On the multivariate transfinite diameter, *Annales Polonici Mathematici* 72 (1999), 285-305.
- [2] J.-B. Bost, H. Gillet and C. Soulé, Heights of projective varieties and positive Green's forms, *Journal of the American Mathematical Society* 7 (1994), 903-1027.
- [3] E. Bedford and B. A. Taylor, A new capacity for plurisubharmonic functions, *Acta Mathematica* 149 (1982), 1-40.
- [4] T. Chinburg, Capacity theory on varieties, *Compositio Math* 80 (1991), 71-84.
- [5] T. Chinburg, C. F. Lau and R. Rumely, Capacity theory and arithmetic intersection theory, *Duke Mathematical Journal* 117 (2003), 229-285.
- [6] H. Gillet and C. Soulé, Arithmetic intersection theory, *IHES Publ Math.* 72 (1990), 94-174.
- [7] H. Gillet and C. Soulé, Characteristic classes for algebraic vector bundles with Hermitian metric, I, II, *Ann. of Math. (2)* 131 (1990), 13-203, 205-238.
- [8] H. Gillet and C. Soulé, An arithmetic Riemann-Roch Theorem, *Invent. Math.* 110 (1992), 473-543.
- [9] M. Klimek, *Pluripotential theory*, Oxford, 1991.
- [10] N. Levenberg and B. A. Taylor, Comparison of capacities in $\mathbb{C}^N$, in: *Complex Analysis (Toulouse, 1983)*, 162-172, *SLNM* 1094, Springer-Verlag, Berlin, 1984.
- [11] R. Richburg, Stetige streng pseudokonvexe Funktionen, *Math. Ann* 175 (1968), 257-286.
- [12] R. Rumely, Capacity theory on algebraic curves, *Lecture Notes in Mathematics* 1378, Springer-Verlag, Berlin-Heidelberg-New York, 1989.
- [13] R. Rumely and C. F. Lau, Arithmetic capacities on $\mathbb{P}^n$, *Math. Zeit.* 215 (1994) 533-560.
- [14] R. Rumely, C. F. Lau, and R. Varley, Existence of the Sectional Capacity, *AMS Memoires* 690 (vol. 145), American Mathematical Society, Providence, R.I., 2000.
- [15] V. P. Sheĭnov, Invariant form of Pólya's inequalities, *Siberian Math. J.* 14 (1973), 138-145.
- [16] V. Zaharjuta, Transfinite diameter, Chebyshev constants, and capacity for compacta in $\mathbb{C}^N$, *Math. USSR Sbornik* vol 25 (1975), 350-364.
- [17] V. Zaharjuta, Extremal plurisubharmonic functions, orthogonal polynomials and Bernstein-Walsh theorem for analytic functions of several variables, *Annales Polonici Mathematici* 33 (1976), 137-148.

ROBERT RUMELY, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GEORGIA, ATHENS, GEORGIA 30602, USA

E-mail address: rr@math.uga.edu